



W&C 21 July 2000

Results from DONUT*: First Direct Evidence of ν_τ

The primary goal is to directly observe the charged-current interactions of the tau-neutrino.

The experiment locates and identifies tau-lepton decays using an emulsion target and spectrometer.

Byron Lundberg FERMILAB
for the DONUT collaboration

**Direct Observation of Nu Tau*



W&C 21 July 2000

DONUT Collaboration

Aichi Univ. of Education

K. Kodama, N. Ushida

Kobe University

S. Aoki, T. Hara

Nagoya University

N. Hashizume, K. Hoshino, H. Inuma, K. Ito,
M. Kobayashi, M. Miyanishi, M. Komatsu,
M. Nakamura, K. Nakajima, T. Nakano, K. Niwa,
N. Nonaka, K. Okada, T. Yamamori

Univ. of California/Davis

P. Yager

Fermilab

B. Baller, D. Boehnlein, W. Freeman,
B. Lundberg, J. Morfin, R. Rameika

Kansas State Univ.

P. Berghaus, M. Kubansteven, N.W. Reay,
R. Sidwell, N. Stanton, S. Yoshida

Univ. of Minnesota

D. Ciampa, C. Erickson, K. Heller, R. Rusack,
R. Schwienhorst, J. Sielaff, J. Trammell, J. Wilcox

Univ. of Pittsburgh

T. Akdogan, V. Paolone

Univ. of South Carolina

A. Kulik, C. Rosenfeld

Tufts University

T. Kafka, W. Oliver, J. Schneps, T. Patzak

Univ. of Athens

C. Andreopoulos, G. Tzanakos, N. Saoulidou

Gyeongsang University

J.S. Song, I.G. Park, S.H. Chung

Kon-kuk University

J.T. Rhee



W&C 21 July 2000

Outline

OVERVIEW of EXPERIMENT

Digression 1: Brief Review of n_t

NEUTRINO BEAM

The INTERACTION sample of 203 EVENTS

Digression 2: Modern Emulsion Primer

DECAY SEARCH

BACKGROUNDS

The SIGNAL

CONCLUSION


Essential quantities



W&C 21 July 2000

Overview

- Use 800 GeV protons \rightarrow beam dump
- Emulsion 36m from dump
- Magnetic/passive shield protects emulsion
- ν flux < 0.001 of NuTeV
- Search for ν_τ interaction by topology: *kinks*





W&C 21 July 2000

Digression 1: The ν_τ Story

1975 - τ lepton discovered ; ν_τ assumed to exist

1980's - CLEO, ARGUS, LEP τ decays: nothing unusual

1986 - E531 : no direct couplings to $\nu_{e,\mu}$

1991 - LEP demonstrates 3.00 ν 's

1998 - Evidence of ν osc. $\nu_\mu \leftrightarrow \nu_\tau$



W&C 21 July 2000

ν_τ Properties

- the usual SM numbers
- mass $< 16 \text{ MeV}/c^2$ (direct)
- dipole moments: $\mu_{\nu_\tau} < 4.2 \times 10^{-7} \mu_B$ (preliminary)



W&C 21 July 2000


τ Decays

- Topological BR $\tau \rightarrow$ one charged particle:

86%

- $\tau \rightarrow e \nu \nu$ 18%

- $\tau \rightarrow \mu \nu \nu$ 18%



- $c\tau_\tau = 87 \mu\text{m} \Rightarrow 2.3\text{mm mean decay length}$
in DONUT



W&C 21 July 2000

Prompt Neutrino Beam

E-872

*Making ν_τ interactions
from protons*







Diagram illustrating the beamline and detector setup for the Muon and Neutrino experiment. The beamline starts with 800 GeV protons entering a Beam Dump, then passes through a SELMA Magnet, a MuSweep2 Magnet, and a Passive Steel Shield. It then enters a Spectrometer, which consists of an Emulsion v Target, followed by muon and neutrino detection regions. The diagram shows the paths of muons (μ^+ and μ^-) and neutrinos (ν) produced in the target. A scale bar indicates 1m and 5m.

- Emulsion Target 36m from beam dump
- Muon rate $\sim 2 \times 10^4$ per 10^{13} *pot* in target area



W&C 21 July 2000


Spectrometer





W&C 21 July 2000


Emulsion Target





W&C 21 July 2000

Emulsion Target / Vertex Detector



- Four target stations
- 260 kg total mass
- Interleaved with sci-fi
- Fibers \rightarrow vtx prediction
- Total 7 modules exposed
- Modules $\sim 2-3 X_0$ each
- $\sim 0.2 - 0.3 \lambda_{\text{int}}$ each



W&C 21 July 2000

Installation of Emulsion Module



April 1997



W&C 21 July 2000

ν Flux / Interactions

Prompt ν beam \Rightarrow number $\nu_e \sim$ number ν_μ

Primary ν_τ source $D_S \rightarrow \nu_\tau \tau \rightarrow \bar{\nu}_\tau X$

$\nu_e + \nu_\mu$	$2.5 \times 10^{-4} / (pot \ m^2)$	52 GeV
ν_τ	$2.1 \times 10^{-5} / (pot \ m^2)$	54 GeV

Total *protons on target* = 3.6×10^{17}

Calculated number of interactions = 1100 (ν_μ , ν_e , ν_τ)

Data taken from April to September 1997



W&C 21 July 2000

ν_τ Sources

- 1. Primary Source: $D_S^+ \rightarrow \nu \tau^+ \rightarrow X \bar{\nu}$


$$BR(D_S^\pm \rightarrow \tau^\pm \nu_\tau) = \left(\frac{G_F^2}{8\pi} \tau_{D_S} m_{D_S} \right) \cdot f_{D_S}^2 |V_{cs}|^2 m_\tau^2 \left(1 - \frac{m_\tau^2}{m_{D_S}^2} \right)^2 = 6.1 \pm 1.0\%$$

- 2. $D^+ \rightarrow \nu \tau^+ \rightarrow X \nu$ (rate 5% of D_S)
- 3. D_S from secondary interactions in dump (rate 8% of D_S)
- 4. $B \rightarrow \tau X$ (rate 1.3% of D_S)



W&C 21 July 2000

How many ν_τ ?



Expected number of
 ν_τ interactions:

4.8% of total

Uncertainties:


D_S production $\pm 20\%$

$f_{D_S} \Rightarrow \text{BR}$ $\pm 15\%$



W&C 21 July 2000

Data Set





W&C 21 July 2000

Data Sample : 203 events


203 events:

- Primary located in emulsion
- Emulsion data well calibrated
- Decay search performed
- Systematic studies *nearly* complete



W&C 21 July 2000

Interaction Vertex Distribution



Predicted interaction
vertices projected in
 x - y plane


Black dots : 203 set

Red dots : 901 predictions



W&C 21 July 2000

ν_μ CC interaction








W&C 21 July 2000

ν_e CC interaction

U View



V View



DONUT

Fermilab E-872

Run= 3250 Event= 470

Electron CC

Station 4

Bulk Emulsion



W&C 21 July 2000

203 set : ν_μ CC Events


75 ν_μ CC interactions
with μ ID'd

47 μ^-

28 μ^+

$$\frac{N(-)}{N(+)} = 1.7$$


expect 1.8





W&C 21 July 2000

203 set: Composition



86 ± 11 ν_μ CC interactions
(acceptance corrected)

Calorimeter energy spectrum:
 79 ± 14 ν_e CC interactions

$\Rightarrow \nu_e \sim \nu_\mu$ in no. interactions

$\Rightarrow \nu_\mu$ non-prompt $< 0.3 \times$ prompt

Recall ν_τ rate 4.8% $\Rightarrow 10 \pm 3$ events



W&C 21 July 2000


Digression 2: Modern Emulsion

- Target design - ECC type vs bulk type
- Digitized emulsion: Analysis flow
- Scanning stations at Nagoya University
- Performance: efficiency, resolution
- Momentum determination using scattering
 - Essential for τ decay analysis



W&C 21 July 2000

Emulsion Plates




AgBr suspended
in a gel is coated
on plastic sheets.

- Fuji ET7C
- Grain size $0.7 \pm 0.2 \mu\text{m}$
- 29 ± 2 grains per $100 \mu\text{m}$ for m.i. track
- *Information capacity* (i.e. $50 \times 50 \times 6 \text{ cm}^3$) :
 $1 \times 10^5 \text{ tracks/cm}^2$ @ $3000 \text{ grains/cm} \Rightarrow$
 $10^{12} \text{ grains} \Rightarrow \sim \text{Terabytes of data}$




W&C 21 July 2000

Target Designs



"ECC 800"



"ECC 200"




"Bulk"

- 3 target types
- Bulk 95% emulsion
- ECC 5% emulsion
- ECC for OPERA



W&C 21 July 2000

Digital Emulsion Analysis





W&C 21 July 2000

Digital Emulsion Analysis: Scanning Stations






W&C 21 July 2000

Emulsion Scan Volume

Decay Search



- Emulsion data digitized, stored on disk
- Vertex, decay search similar to electronic detectors




Emulsion Scanning Efficiency:

1^{ry} Location

W&C 21 July 2000

- Location efficiency factored into two independent parts:
 - Expect uniform(flat) z-vertex distribution within module. But ... Easier to find interactions on downstream edge of module:






W&C 21 July 2000

Emulsion Scanning Efficiency: 1^{ry} Location (*cont.*)

→ Higher multiplicity events are easier to find. Some bias against low multiplicity interactions.





W&C 21 July 2000

Emulsion Scanning Efficiency:

1^{ry} Location

- From the shape of located z-vertex distribution and comparing the expected 1^{ry} charged track multiplicity to the data one can extract the following 1^{ry} location efficiencies for different ν interaction types*:


<i>ν interaction type</i>	<i>Location Efficiency</i>
ν_{cc} (e, μ 's only)	52 \pm 10%
ν_{nc} (all types)	46 \pm 9%
ν_{cc} + Charm	52 \pm 10%
ν_{τ} cc	55 \pm 11%

*Overall location efficiency from data: (located/attempted) = 52 \pm 6%



W&C 21 July 2000

Emulsion Performance: Plate Efficiency



**Tracking Efficiency
for 2632 plates**


Track $\theta < 0.1$ rad

**Only 24 plates with
connection efficiency
below 90%**



W&C 21 July 2000

Emulsion Performance: Spatial Resolution






The emulsion track
residuals

Average : 0.29 μm



W&C 21 July 2000

Emulsion Performance: Spatial Resolution




Emulsion data calibrated to $0.3 \mu\text{m}$ in transverse coordinates ($0.4 \mu\text{m}$ for bulk emulsion)

Typical vertex precision (C5) $0.5 \mu\text{m}$ transverse, and $15 \mu\text{m}$ along beam




W&C 21 July 2000

Emulsion Performance: Vertex Precision



Distance of approach
of emulsion tracks to
the primary vertex
with $\theta < 0.1$ rad




$rms = 0.8 \mu\text{m}$



W&C 21 July 2000

Momentum Measurement using Multiple Scattering in Emulsion Targets




$$\frac{\Delta\vartheta^{rms}}{\vartheta^{rms}} \propto \frac{1}{\sqrt{n}} \quad n : \text{number of measurements}$$

For example: 10 GeV/c has *rms* deflection of 0.3μm



W&C 21 July 2000

Momentum from Scattering: Test Beam Results




**Momentum Measured
using Multiple Scattering**

Beam Test

4 GeV/c π^-


Stack of 29 bulk plates





W&C 21 July 2000

Momentum from Multiple Scattering: Calibration in DONUT




$rms = 35\%$



W&C 21 July 2000

Momentum from Multiple Scattering: Electron Energy Loss



Electron ID / energy loss

Scattering method applied
to e^\pm is usually a lower
limit to p_e



W&C 21 July 2000

Intermission

- The achieved emulsion performance exceeds our initial expectations
- Located interactions have been verified in the SFT
- The 203 data set is fully calibrated
- We can measure momenta with emulsion data, reliably to 20 GeV/c
 - *Important for decay search*



W&C 21 July 2000


Decay Search

- Strategies: “Long” kinks
“Short” kinks
- Cuts on kink data for τ search
- Decay search efficiency estimate




W&C 21 July 2000

C1 Decay Search



1. *Long Decays*

- parent measured
- kink resolved
- $\tau \Rightarrow$ no 1^{ry} lepton
- $\sim 75\%$




2. *Short Decays*

- IP wrt 1^{ry} vertex
- only daughter meas.
- daughter seen in spect.
- $\sim 25\%$



W&C 21 July 2000

Neutral Decay Search



3. *Neutral* search

- *charm* only
- daughters in spect.



W&C 21 July 2000

Decay Search: Kink Selection


<i>Cut</i>	kinks	$\% \tau$ remaining*
none	7642	100
0.01 < kink angle < 0.250	450	78
Parent angle < 0.20 rad	280	90
IP (parent) < 5 μ m	142	>99
IP (@ kink) < 5 μ m	65	>99
Flight length < 5mm	42	78
C1 (kink)	42	86
“Long” decay		76
Total		72

*not cumulative



W&C 21 July 2000

Decay Search: Kink Finding Efficiency



Kink finding efficiency
as a function of p_T

for $p_T > 250 \text{ MeV/c}$:

$$\epsilon_{\text{kink}}(\tau) = 65\%$$


$$\epsilon_{\text{kink}}(D^\pm) = 56\%$$




W&C 21 July 2000

Kinks: Classification


- Randomly associated tracks
 - e.g. Primary track + stale muon track




- *Charm* background



- Interactions (scattering)





- Tau signal





W&C 21 July 2000

Backgrounds: Random Association




Procedure: “Break” a track from the primary into two pieces, then find the distance between the two, refit tracks

$$rms = 1 \mu\text{m}$$



W&C 21 July 2000

Backgrounds: Random Association



For each event, *all* tracks starting within 1 plate of the vertex are shown

The shape is characteristic of uncorrelated, uniform distribution:


⇒ 80 tracks/mm²/event

Est. random kinks < 10⁻³



W&C 21 July 2000

Backgrounds: Random Association




Same set of track as in
previous slide, but showing
only IP < 20 μm

Only 6 tracks within 5 μm



W&C 21 July 2000



Background : *charm* Production

1. assume $\nu_\mu \sim \nu_e$	168 ± 18	no. CC int.	
2. <i>charm</i> production	0.081 ± 0.008	13.6 ± 1.3	
3. charged charm	0.47 ± 0.05	6.4 ± 0.9	
4. “Long” decay & $< 5\text{mm}$	0.66 ± 0.06	4.2 ± 0.7	
5. kink detection effic.	0.56 ± 0.06	2.4 ± 0.5	
6. <i>charm</i> \rightarrow kink (C1)	0.38 ± 0.02		
$\Pi_{1..6}$	0.9 ± 0.2 events	[1.5 (inflate all 1σ)]	

Expect 2.4 charged *charm* decays : found 2




Expect 0.9 *charm* kink decays : found 1



W&C 21 July 2000

charm Production



Neutrino-produced
charm data


CCFR
E531

~~For DONUT
charm to total avg:
0.068 ± 0.004~~

Update:
NuTeV preliminary
results: 0.078 ± 0.007



W&C 21 July 2000



Background: *charm* summary

We estimated 0.9 ± 0.2 *charm* events with *kink* :

This is a background to τ *iff* the *lepton* is not identified as such

- μ ID prop tubes cover 82% of acceptance \times 96% eff. = 79%
- e ID require $>2X_0$ for emulsion tag or $>3X_0$ FT : 75%


\therefore *Total charm background* : $0.23 \times 0.9 = 0.21 \pm 0.04$ events



W&C 21 July 2000

τ Background : *Interactions*

“NC interactions” + hadron scatter = τ background



- Rapidly decreasing with p_T
- Depends on total path length
- *Short* decays is separate analysis

Estimated background using
GEANT calculation :


0.20 ± 0.06 events
for $p_T > 250$ MeV/c



W&C 21 July 2000

Background: Interactions

Distribution of Kinks



All kinks from analysis
of the 203 set with:

$$0.01 < \theta(\text{ kink}) < 0.25;$$

$$\theta(\text{parent}) < 0.20;$$

$$\text{IP}(\text{parent}) < 5 \mu\text{m};$$

$$\text{IP}(@\text{kink}) < 5 \mu\text{m}$$


65 tracks out of 654



W&C 21 July 2000

Background: Interactions

Distribution of Kinks



Kinks for which p_T is computed using scattering

It does not include all kinks from kink list


Recall: only tracks in “NC-like” events are part of τ background (~39%)



W&C 21 July 2000

Background+Signal: Interactions

Distribution of Kinks




Same plot as previous slide,
but *color enhanced* !



W&C 21 July 2000

3333 17665 Magnified






W&C 21 July 2000

Background+Signal: Interactions

Distribution of Kinks



What if *each* signal event
is a $+1\sigma$ fluctuation ?

Lower momentum for all
tau events by 1σ

Signal still separated!



W&C 21 July 2000

Summary: τ candidates

from 203 set

- 4 events from *Long* decay search;

$$\text{expect } (203)(0.048)(0.86)(0.76)(0.65) = 4.1 \pm 1.4$$

τ frac C1 decay *Long* kink eff

Background Analysis:

Charm : 0.21 ± 0.04 events

Hadronic interactions : 0.20 ± 0.06 events

Sum Background: 0.41 ± 0.15 events

Fluctuation bkg \rightarrow signal : Poisson prob. 8×10^{-4}



W&C 21 July 2000

Next steps

- “Short” Decay Analysis to be completed
- Complete detailed characterization of data set
- The data set *can* be doubled ...



W&C 21 July 2000

In Conclusion

- “Long” Decay Search kink list only 65 events (after cuts)
- Randomly assoc. kinks $<10^{-3}$ for entire 203 event set
- *charm* background for τ calculated at 0.21 ± 0.04 events
- Interaction background calc. at 0.20 ± 0.06 events ($p_T > 250$)
 - Kink data supports MC calc.
- Signal events far from kink background in p_T

⇒ We conclude that these events are the evidence for observing the process:

$$\nu_\tau + N \rightarrow \tau + X$$



W&C 21 July 2000

Acknowledgements

DONUT would like to thank the Fermilab support staff for helping make our little experiment a big success